# **VACCINE POINTS**

"Vaccine Points" is a bi-monthly communication distributed by the Hawaii Department of Health to all participating Vaccines for Children (VFC) providers.

Test your vaccine knowledge.

## **Question:**

Which of the following vaccines, if any, is/are recommended by the ACIP for administration to a 6 month old child? Select all that apply.

- a) Menactra<sup>®</sup> (child moving to an area where meningococcal disease is hyperendemic)
- b) Varivax® (sibling diagnosed with chickenpox this morning)
- Rotarix® (child received Rotarix® at ages 2 and 4 months)
- d) Hiberix® (child received doses of ActHIB® at aged 2 and 4 months)
- e) All of the above
- None of the above

#### askexperts/experts hib.asp

Immunization Action Coalition Ask the Experts: Haemophilus influenzae type b (Hib) http://www.immunize.org/

(Hib) Vaccination Series, April 29, 2016, p. 418. https://www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6516a3.pdf MMWR Food and Drug Administration Approval for Use of Hiberix as a 3-Dose Primary Haemophilus influenza Type b

Committee on Immunization Practices (ACIP), February 6, 2009, p. 17. http://www.cdc.gov/mmwr/PDF/rr/rr5802.pdf MMWR Prevention of Rotavirus Gastroenteritis Among Infants and Children Recommendations of the Advisory

June 22, 2007, p. 23. https://www.cdc.gov/mmwr/pdf/rr/rr5604.pdf

MWWR Prevention of Varicella: Recommendations of the Advisory committee on Immunization Practices (ACIP),

Immunization Action Coalition Ask the Experts: Varicella (chickenpox) http://www.immunize.org/askexperts/

### http://www.cdc.gov/mmwr/pdf/wk/mm6324.pdf

Disease: Recommendations of the Advisory Committee on Immunization Practices, 2013, page 527. MMWR Use of MenACWY-CRM Vaccine in Children Aged 2 Through 23 Months at Increased Risk for Meningococcal 2onces:

at ages 2, 4, and 6 months. All of the licensed Hib-containing vaccines may be used interchangeably. January 2016, the FDA approved the expanded use of Hiberix® for a 3-dose infant primary vaccination series Hiberix® was first licensed in 2009 for use as a booster dose in children aged 15 months through 4 years. In

Rotarix® is administered in a 2-dose series, with doses administered at ages 2 and 4 months.

Varicella vaccine is neither approved nor recommended for children younger than 12 months of age.

Menactra<sup>®</sup> is licensed for administration to children aged 9 months and older.

